

## HOW ‘PASTORAL’ IS PASTORALISM? DIETARY DIVERSITY IN BRONZE AGE COMMUNITIES IN THE CENTRAL KAZAKHSTAN STEPPES\*

E. LIGHTFOOT,<sup>1</sup>† G. MOTUZAITE-MATUZEVICIUTE,<sup>2</sup> T. C. O’CONNELL,<sup>3</sup>  
I. A. KUKUSHKIN,<sup>4</sup> V. LOMAN,<sup>4</sup> V. VARFOLOMEEV,<sup>4</sup> X. LIU<sup>1</sup> and M. K. JONES<sup>3</sup>

<sup>1</sup>McDonald Institute for Archaeological Research, University of Cambridge, Downing Street, Cambridge CB2 3ER, UK

<sup>2</sup>History Faculty/Department of Archaeology, Vilnius University, Universiteto 7, 01513 Vilnius, Lithuania

<sup>3</sup>Department of Archaeology and Anthropology, University of Cambridge, Downing Street, Cambridge CB2 3DZ, UK

<sup>4</sup>Saryarka Archaeological Institute, Karaganda State University named after EA Buketov, 28 University Street, Karaganda 100028, Kazakhstan

*Steppe communities have traditionally been viewed as pastoralist groups with similar herd-based economies. Recent scholarship, however, warns against assumptions of homogeneity and new scientific techniques are providing a more nuanced approach to steppe archaeology, with increasing indications of diversity. This recent evidence further suggests that considering these communities as primarily pastoralist may hide a variety of subsistence strategies, such as fishing and cultivation. Here, we consider direct evidence for diet (in the form of stable isotope analysis) from Bronze Age communities from central Kazakhstan, in the semi-arid steppe zone. We find that the diversity recently suggested for communities across the steppe zone can be found within sites in the Karaganda region. This suggests that individuals exercised choice in their dietary habits that led to dietary differences large enough to be detectable isotopically. The results also highlight the inclusion of fish in the diet of these ‘pastoral’ populations, with indications that some individuals in the Final Bronze Age consumed notable amounts of millet. This shows that these ‘pastoralist’ economies also engaged in fishing throughout the Bronze Age, with millet cultivation becoming increasingly important in the Final Bronze Age. As such, our understanding of what it means, in this context, to be a pastoralist requires further consideration.*

KEYWORDS: PALAEODIET, CARBON, NITROGEN, STEPPE

### INTRODUCTION

Stretching from Hungary to eastern Russia, the steppe ecological zone has played a vital part in past world events, and understanding the archaeology of the steppe is crucial to a full understanding of world prehistory. A central theme in the archaeology of the steppe has been the idea of Bronze Age pastoralists, and the role their ranging lifestyle played in the spread of a variety of attributes, including horse husbandry, metal technology and Indo-European language (cf., Gimbutas 1973; Renfrew 1987; Anthony 2007). However, the cultures populating the steppes were, until relatively recently, described by large-scale models that lacked good connections to local adaptations, socio-economic change and the mosaic of environments that characterize the steppes (Hanks 2010: cf., Frachetti 2006; Bendrey 2011). Nevertheless, within the past 20 years international interest and research into steppe archaeology has significantly increased, and new archaeological techniques are allowing a more nuanced understanding of steppe archaeology.

\*Received 3 March 2014; accepted 15 May 2014

†Corresponding author: email elfl2@cam.ac.uk

© 2014 University of Oxford

Several recent projects have focused on the Bronze Age (such as Frachetti and colleagues in South Eastern Kazakhstan: Frachetti 2002, 2006), a particularly dynamic period that saw the emergence of large-scale patterns of movement, interaction and trade between regional communities and polities, as well as key technological developments such as spoke-wheeled chariots (Hanks 2010). These projects have tended to highlight the diversity of human adaptation to the steppe environments, and provide some indications that viewing these communities as simply pastoralist may limit our awareness of other aspects of their subsistence. For example, Bendrey (2011) found that herd composition varied across the steppes influenced by regional climatic, topographical and ecological conditions, while Frachetti and colleagues (2010) have found evidence for domesticated plants, indicating that terrestrial animals were not the sole focus of food production.

Here, we use stable carbon and nitrogen isotope analysis to consider the diversity of human diets in Bronze Age central Kazakhstan and to assess critically what it meant, in dietary terms, to be a 'pastoralist' in this semi-arid steppe zone. Through the use of this technique we will provide independent and direct dietary evidence at the site, regional and, through comparison to published research, international levels. We will also consider the consumption of resources other than domesticated animals, such as fish and cultivated plants (specifically C<sub>4</sub> plants), which may be absent from the archaeological record of these communities.

## BACKGROUND

### *Kazakhstan archaeology and evidence for diet*

The Bronze Age in central Kazakhstan is divided into a number of cultures represented in this study: the Late Bronze Age Andronovo culture, divided into Alakul and Fedorovo sub-types (mid-second millennium BC); and the Final Bronze Age Begazy–Dandybai culture (1300–900 BC: Hanks *et al.* 2007; Koryakova and Epimakhov 2007; Svyatko *et al.* 2009). All of these cultures are thought to have subsistence practices focusing on pastoralism. Zooarchaeological remains are increasingly studied, and a limited amount of lipid residue analysis has been carried out, but flotation for archaeobotanical evidence remains uncommon.

Throughout the Bronze Age, the animal bone evidence indicates that the economy was based upon caprids, cattle and horses, with limited use of large wild animals (Vadetskaya 1975, as cited in Frachetti 2002; Kuzmina 2003 and references therein; Outram *et al.* 2012). By the Andronovo period there is also evidence for pigs and dogs. Sieving has been limited, but was carried out at Andronovo Temirkash (sampled in this study), indicating that bird and fish bones can be found but are present in very low proportions (1.1% and 1.5%, respectively; Outram *et al.* 2012). Lipid residue and herd structure analyses from the Andronovo site of Lisakovsk (in the forest-steppe zone of north-west Kazakhstan) and the Begazy–Dandybai proto-city Kent (analysed in this study) indicates that, at these sites at least, ruminant dairy products were utilized, particularly from cows (Outram *et al.* 2012).

As noted above, archaeobotanical studies are rare; however, Bronze Age plant remains have been recovered from Begash in south-east Kazakhstan and included two types of domesticated grains, broomcorn millet (*Panicum miliaceum*) and free-threshing wheat, directly dated to 2460–2150 cal BC (Frachetti *et al.* 2010; Spengler *et al.* 2013, 2014). While this indicates that domesticated crops may be present, they were largely absent from domestic contexts, which suggests ritual rather than subsistence use. Spengler and colleagues (2014) also report a small number of wheat grains from Tasbas in south-eastern Kazakhstan, dated using charcoal to

2840–2500 cal BC, while Late Bronze Age flotation samples from Tasbas contained much higher seed densities, including naked barley, broomcorn millet and peas, with barley grains directly dated to 1450–1250 cal BC. The only other early finds of wheat and millet are from the Middle Bronze Age sites of Arkaim and Alandskoe (c.2200–1800 BC) in the trans-Ural region (Gayduchenko 2002); however, the report lacks species identification, direct chronology and morphological information (Frachetti *et al.* 2010). In the Iron Age, there is well-dated evidence for domesticated plant use in the form of millet, wheat and barley phytoliths from Tuzusai and Tseganka 8 in southeastern Kazakhstan (Chang *et al.* 2003).

### *Stable isotope analysis*

Stable isotope analysis provides individual, quantitative evidence about past diets. As the body is made from the food and drink consumed during life, chemical differences in diets are transmitted through the food chain and recorded in bone collagen. The technique is relatively insensitive to minor dietary components, and it is estimated that 20% of the protein consumption must derive from an isotopically distinct source in order for it to be detectable isotopically (Hedges 2003). This means that when isotopic differences are found, they reflect real differences during life.

Stable carbon isotope ratios ( $\delta^{13}\text{C}$ ) can be used to distinguish between diets based upon two different types of plant ( $\text{C}_3$  and  $\text{C}_4$ ), which follow different photosynthetic pathways (Vogel and Van der Merwe 1977). Most economically important plants, including wheat, barley, rice and potatoes, are  $\text{C}_3$ , but maize, millet, sorghum and sugar cane use the  $\text{C}_4$  photosynthetic pathway. At this time, millet is the only  $\text{C}_4$  staple crop that would have been available to people living in central Kazakhstan. However, it is not necessarily the only  $\text{C}_4$  plant entering the human food chain. The steppe zone includes numerous  $\text{C}_4$  plants present in relatively small proportions (Pyankov *et al.* 2000; Toderich *et al.* 2004). While it is difficult to imagine a scenario whereby direct consumption of these plants by humans would notably influence their  $\delta^{13}\text{C}$  values, it is possible that animals ate isotopically detectable amounts of wild  $\text{C}_4$  plants. Stable carbon isotope ratios can also be used to distinguish between marine and terrestrial food chains (Schoeninger and De Niro 1984), but marine foods are unlikely to have been available to people in prehistoric central Kazakhstan.

Stable nitrogen isotope ratios ( $\delta^{15}\text{N}$ ) provide an indication of the amount of animal protein consumed by an individual, as the  $\delta^{15}\text{N}$  value increases by 3–5‰ with each step in a trophic chain, although the mechanism that causes this is not clearly understood (O’Connell and Hedges 1999; Bocherens and Drucker 2003; Hedges and Reynard 2007). Aquatic ecosystems tend to have long food chains, which leads to higher  $\delta^{15}\text{N}$  values than in terrestrial food chains (Schoeninger and De Niro 1984).

In this study, bone collagen (protein) was analysed, which reflects diet over the majority of adult life, and small dietary inputs or changes in diet may not be detectable (Hedges *et al.* 2007). As nitrogen is only present in dietary protein, the nitrogen isotope values reflect only the protein component of the diet. Carbon in bone collagen is mainly taken from the protein part of the diet, although some carbon may reflect other dietary fractions (Ambrose and Norr 1993; Tieszen and Fagre 1993; Howland *et al.* 2003; Jim *et al.* 2006). Infants tend to have higher  $\delta^{15}\text{N}$  values than adults due to a trophic level effect associated with breastfeeding (Fuller *et al.* 2006). Pathological conditions and starvation can lead to increases in body  $\delta^{15}\text{N}$  values (Katzenberg and Lovell 1999; Mekota *et al.* 2006). When conducting stable isotope analyses on human populations, it is beneficial to sample contemporaneous fauna, as the isotopic ratios at the base of the food chain vary through space and time (Stevens and Hedges 2004).

## MATERIALS AND METHODOLOGY

Eighty-eight human bone samples and 76 animal samples (horse, ovicaprid, cattle and dog) were taken from sites in central Kazakhstan (Fig. 1 and Table 1). Full sample details and isotope results are presented in Supplementary Appendices 1 and 2. The human samples span the Late to Final Bronze Age, with three samples dated to the Early Iron Age. The samples are dated on the basis of associated material culture (for details, see the references in Table 1) and, in the absence of direct dating, it remains possible that some of the individuals are misdated, particularly as the kurgans could have been reused in later time periods (potentially without any associated material culture to signify this). The majority of the animal samples date to the Andronovo period; that is, the later Bronze Age.

Collagen was extracted following the method described in Privat *et al.* (2005). All collagen samples were analysed in triplicate, using a Costech elemental analyser coupled in continuous-flow mode to a Thermo Finnigan MAT253 mass spectrometer. Carbon and nitrogen stable isotope values are expressed as delta values (e.g.,  $\delta^{13}\text{C}$ ) relative to international standards (VPDB and AIR, respectively) in units of per mille (parts per thousand, ‰; Hoefs 2004). Repeated measurements on international and in-house standards showed that the analytical error was less than <0.2‰ for both carbon and nitrogen.

Measured collagen is deemed to be of good quality if it fulfils the following criteria: an atomic C : N ratio of 2.9 to 3.6 (De Niro 1985); a 'collagen' yield of >1% by mass; final carbon yields of >13%; and final nitrogen yields of >4.8% (Ambrose 1990). Three human and six animal samples failed to yield reliable collagen. All of the remaining samples yielded reliable collagen: the C : N ratios of all samples fell within a range of 3.1 to 3.4; and all collagen yields were greater than 1%.

Statistical analyses were performed using SPSS 21.0 for Mac. Samples were tested for normality using histograms, and Kolmogorov–Smirnov and Shapiro–Wilk tests, and for equality of variance using Levene's tests. The parametric data were investigated using independent samples *t*-tests. The non-parametric tests employed were Kruskal–Wallis tests. All statistical data are given in Table 2. Outliers are identified as samples that lie more than 1.5 times the interquartile range (IQR) below quartile 1 (Q1) or above quartile 3 (Q3). The minimum sample size for statistical analysis was five.

## RESULTS

*Animal results*

The animal results have been grouped into herbivores and carnivores and are shown in Figure 2. The herbivore values (horse, ovicaprid and cattle) have a mean  $\delta^{13}\text{C}$  value of  $-19.2\text{‰}$  ( $\pm 0.7\text{‰}$ ) and  $\delta^{15}\text{N}$  of  $7.5\text{‰}$  ( $\pm 1.6\text{‰}$ ), and range from  $-20.6$  to  $-17.9\text{‰}$  in  $\delta^{13}\text{C}$  and from  $4.8\text{‰}$  to  $13.3\text{‰}$  in  $\delta^{15}\text{N}$ . While there are no statistically identified outliers in  $\delta^{13}\text{C}$ , two of the caprid samples (KZF44 and KZF56, from Temirkash and Tashik, respectively) have relatively high  $\delta^{13}\text{C}$  values (both  $\delta^{13}\text{C} = -17.9\text{‰}$ ), which may suggest a small  $\text{C}_4$  component to their diets or the consumption of ( $\text{C}_3$ ) plants with high  $\delta^{13}\text{C}$  values due to aridity (O'Leary 1988; Farquhar *et al.* 1989; Hartman 2012). Three samples are identified as statistical outliers in  $\delta^{15}\text{N}$  (lying more than 1.5 times the IQR above Q3; KZF46, KZF47 and KZF50), with values greater than  $12.5\text{‰}$ . These individuals could be: young animals exhibiting a weaning effect; subject to extreme stress, such as starvation (Mekota *et al.* 2006); imported from an area with very high  $\delta^{15}\text{N}$  values; or




Figure 1 A map of the analysed sites.

Table 1 A brief description of the analysed sites

Site	Culture/period	Date	Comments	Number of samples	Reference(s)
Tegiszhol (Тегисжол)	Andronovo (Atasu–Alakul, Nura–Fedorovo, Begazy–Dandybai) and Tasmola/Bronze Age and Early Iron Age	1800–700 BC	Kurgan cemetery associated with Temirkash; different dates for different kurgans	29 humans (2 failed) 4 animals	Outram <i>et al.</i> (2011, 2012), Varfolomeev (2007a,b, 2011b)
Temirkash (Темиркаш)	Andronovo/Middle and Late Bronze Age Могильник Темиркаш (cemetery)/Final Bronze Age Поселение Темиркаш/Middle Bronze Age Andronovo–Fedorovo/Final Bronze Age	Mid-second millennium BC and beginning of first millennium BC	Settlement associated with Tegiszhol; located north-west of Karaganda, in semi-arid steppe	2 humans (могильняк) 40 animals (поселение)	Outram <i>et al.</i> (2011, 2012), Varfolomeev (2007a)
Nurataldy (Нураталды)	Early Andronovo – early Alakul	18th to 16th centuries BC	Buried in stone box, with pot of sheep bones	1 human 1 sheep 3 humans	Unpublished
Darıya (Дарья)	Andronovo–Fedorovo				Loman and Kukushkin (2012)
Tashik (Ташик)	Andronovo–Alakul/Middle Bronze Age			15 humans 20 fauna	Unpublished
Aschisu (Ащису)	Early Andronovo – early Alakul	End of third, beginning of second millennium BC	One individual buried with a copper bowl	9 humans 4 fauna	Kukushkin (2007, 2010, 2012), Kukushkin and Loman (2011)

Table 1 (continued)

Site	Culture/period	Date	Comments	Number of samples	Reference(s)
Ауар-берген (аял-берген)	Early Andronovo – early Alakul	18th to 17th centuries BC		1 human	Kukushkin (2006)
Кора-1 (Копка-1)	Andronovo–Alakul		Cemetery	1 human	Unpublished
Ақимбек (Ақимбек)	Andronovo–Fedorovo			5 humans 1 horse	Unpublished
Тасырбай (Тасырбай 2)	Andronovo–Fedorovo			2 humans	Unpublished
Қудуғауауа Сорка-1 (Қудрявая Сорка-1)	Sargarinsko–Alekseevsk/Final Bronze Age			2 humans	Loman and Kukushkin (2009)
Қузылқол (Қызылқол)	Andronovo – early Alakul	18th to 17th centuries BC	Woman buried with seven children	1 sheep 8 humans	Unpublished
Қузыл (Қызыл)	Final Bronze Age	By analogy, 9th to 8th centuries BC		2 humans	Beisenov and Varfolomeev (2008)
Қаратүгі (Қаратүгі)	Sargarinsko–Alekseevsk/Final Bronze Age	By analogy, 9th to 8th centuries BC		5 humans	Unpublished
Жартастас (Жартастас)	Early Iron Age			1 human	Kukushkin and Loman (2004)
Қайрақты (Қайрақты)	Andronovo–Fedorovo		Sacrificial victim	1 human	Loman and Kukushkin (2011)
Кент (Кент)	Begazy–Dandybai/Final Bronze Age	12th to 8th centuries BC	Proto-city	1 human	Outram et al. (2011, 2012), Varfolomeev (2011a, 2013)

Table 2 Statistical data

Comparison	n	$\delta^{13}C$			$\delta^{15}N$				
		Test	Degrees of freedom	Test statistic	P	Test	Degrees of freedom	Test statistic	P
Temirkash versus Tashik animal results	40, 16	t-test	54	(t) 0.041	0.967	t-test	54	(t) -0.455	0.651
Temirkash animal results MBA versus LBA	19, 21	t-test	38	(t) 0.124	0.902	t-test	38	(t) -0.231	0.818
Comparison of human results by site (with $n > 4$ )	26, 15, 9, 5, 8, 5	Kruskal-Wallis	5	(H) 9.584	0.088	ANOVA	5, 62, 67	(F) 3.512	0.007
Comparison of human results by culture ( $n > 4$ )	19, 27, 15, 16	ANOVA	3, 73, 76	(F) 7.382	<0.001	Kruskal-Wallis	3	(H) 4.649	0.199






Figure 2 A scatter plot of the human and animal  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  results.

misidentified (this latter hypothesis is particularly likely for sample KZF47, which was taken from a rib). Further, sample KZF53 has a high  $\delta^{15}\text{N}$  value compared to the remaining herbivores—in this case, the sample was identified as a juvenile at the time of sampling and therefore probably exhibits a weaning effect. These four outliers have been excluded from subsequent analyses. With these outliers excluded, the mean values are  $-19.2\text{‰}$  ( $\pm 0.6\text{‰}$ ) in  $\delta^{13}\text{C}$  and  $7.2\text{‰}$  ( $\pm 1.0\text{‰}$ ) in  $\delta^{15}\text{N}$  (ranging from  $-20.6\text{‰}$  to  $-17.9\text{‰}$  in  $\delta^{13}\text{C}$  and from  $4.8\text{‰}$  to  $8.9\text{‰}$  in  $\delta^{15}\text{N}$ ).

The two carnivore values are notably higher in  $\delta^{15}\text{N}$  than the mean herbivore value ( $\delta^{15}\text{N} = 11.3\text{‰}$  and  $10.0\text{‰}$ ) and have relatively high  $\delta^{13}\text{C}$  values ( $\delta^{13}\text{C} = -18.2\text{‰}$  for both).

Animal samples were available from seven sites, but only two of those sites have a herbivore sample size greater than five: Temirkash ( $n = 39$ ) and Tashik ( $n = 15$ , with two outliers identified above excluded). Statistical analyses show that there is no difference between the sites in either mean  $\delta^{13}\text{C}$  or  $\delta^{15}\text{N}$ .

It is difficult to compare the faunal results through time, as the majority of the samples date to the Andronovo period (see Supplementary Appendix 2). The samples from Temirkash can, however, be divided into Middle and Late Bronze Age. Statistical analyses indicate that there is no difference in either mean  $\delta^{13}\text{C}$  or  $\delta^{15}\text{N}$ .

Although the data are limited, there seems to be little variation in the environmental baseline through space and time in Bronze Age central Kazakhstan. The animal data presented will therefore be combined and used, with caution, as a baseline for all of the human values.

### Human results

The human results are shown in Figure 2 and have a mean  $\delta^{13}\text{C}$  value of  $-18.3\text{‰}$  ( $\pm 0.6\text{‰}$ ) and a mean  $\delta^{15}\text{N}$  value of  $13.3\text{‰}$  ( $\pm 1.3\text{‰}$ ). However, there is one sample with a very low  $\delta^{15}\text{N}$  value

that is statistically identified as an outlier (lying more than 1.5 times the IQR below Q1; KZ006  $\delta^{15}\text{N} = 7.0\text{‰}$ ). This value is more typical of the herbivore than human data in this study. We suggest that this (rib) sample was misidentified and thus we exclude it from further analysis. A second sample also stands out as unusual: sample KZ053 has the highest  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  values of any sample in this study ( $\delta^{13}\text{C} = -15.7\text{‰}$ ,  $\delta^{15}\text{N} = 17.7\text{‰}$ ) and is a statistical outlier in both  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  (lying 1.5 times the IQR above Q3 for both isotopic values). In order to prevent this sample from skewing the means and statistical tests, we exclude it from statistical analyses but include it in the discussion below. With these two samples excluded ( $n = 83$ ), the mean  $\delta^{13}\text{C}$  value is  $-18.3\text{‰}$  ( $\pm 0.6\text{‰}$ , range from  $-19.2\text{‰}$  to  $-16.1\text{‰}$ ) and the mean  $\delta^{15}\text{N}$  value is  $13.3\text{‰}$  ( $\pm 1.0\text{‰}$ , range from  $11.4\text{‰}$  to  $15.7\text{‰}$ , although note that this includes juvenile individuals—when identified juveniles are removed, mean  $\delta^{15}\text{N} = 13.2\text{‰}$ ). When these mean human values are compared to the herbivore data presented above, the mean human–herbivore offset values are  $1.0\text{‰}$  in  $\delta^{13}\text{C}$  and  $6.1\text{‰}$  in  $\delta^{15}\text{N}$ . The herbivore–human difference at Tegiszhol and Temirkash (which are a settlement and an associated cemetery) is  $1.0\text{‰}$  in  $\delta^{13}\text{C}$  and  $5.9\text{‰}$  in  $\delta^{15}\text{N}$  ( $1.1\text{‰}$  and  $6.0\text{‰}$ , respectively, with juveniles removed), while that at Tashik is  $0.8\text{‰}$  in  $\delta^{13}\text{C}$  and  $6.0\text{‰}$  in  $\delta^{15}\text{N}$  ( $0.7\text{‰}$  and  $5.7\text{‰}$ , respectively, with juveniles removed).

Six sites have human sample sizes greater than four: Tegiszhol ( $n = 26$ ), Tashik ( $n = 15$ ), Aschisu ( $n = 9$ ), Akimbek ( $n = 5$ ), Kyzylkol ( $n = 8$ ) and Karatugai ( $n = 5$ ). As can be seen in Figure 3, there is no trend in values with site; rather, there are a range of isotopic values within each site. There are no statistical differences between sites in  $\delta^{13}\text{C}$ , and the only statistical differences in  $\delta^{15}\text{N}$  are that Aschisu has lower values than Akimbek and Kyzylkol.

Finally, the results are compared by culture (and period). As can be seen in Figure 4, in general there is very little difference between periods, with the exception that the Final Bronze Age has more samples with high  $\delta^{13}\text{C}$  values. Statistical tests were run on the Early Alakul ( $n = 19$ ), Alakul ( $n = 27$ ), Fedorovo ( $n = 24$ , excluding sample KZ053 discussed above) and Final Bronze




Figure 3 A scatter plot of the human  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  results, separated by site.




Figure 4 A scatter plot of the human  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  results, separated by culture/period.

Age ( $n = 17$ ) groups. No statistical differences were found in  $\delta^{15}\text{N}$ . In  $\delta^{13}\text{C}$ , the Final Bronze Age was different to all of the other groups, having higher  $\delta^{13}\text{C}$  values.

## DISCUSSION

### Animals

The herbivore  $\delta^{13}\text{C}$  values are typical for Holocene  $\text{C}_3$  environments, showing that the majority of the herbivore diets consisted of  $\text{C}_3$  foodstuffs. There are two caprid results, noted above, with relatively high  $\delta^{13}\text{C}$  values, which may indicate either a small  $\text{C}_4$  input to the diet or more arid conditions at that time. That these two caprids are from different sites indicates that this  $\text{C}_4$  plant(s) and/or aridity effect was present at more than one location. Both samples date to the Andronovo period.

The herbivore  $\delta^{15}\text{N}$  results (mean  $\delta^{15}\text{N} = 7.3\text{‰}$ ) are higher than those found in the Minusinsk basin ( $n = 18$ , mean  $\delta^{15}\text{N} = 5.4\text{‰}$ ; Svyatko *et al.* 2013) and than Eneolithic horses from northern Kazakhstan ( $n = 4$ ,  $\delta^{15}\text{N} = 2.7\text{‰}$ ), but comparable to herbivores from Liskavosk ( $n = 5$ ,  $\delta^{15}\text{N} = 7.5\text{‰}$ ) and Bestamak ( $n = 3$ ,  $\delta^{15}\text{N} = 6.7\text{‰}$ ) in northern Kazakhstan (Ventresca Miller *et al.* 2014), and to those from sites in southern Siberia ( $n = 4$ ,  $\delta^{15}\text{N} = 7.5\text{‰}$ ; Murphy *et al.* 2013). High  $\delta^{15}\text{N}$  values can be associated with manuring (which adds high  $\delta^{15}\text{N}$  biomass to the soil; Bogaard *et al.* 2007), but given the ecological context of the steppes this seems unlikely. Rather, we suggest that aridity is a more likely cause, as aridity has been shown to increase  $\delta^{15}\text{N}$  values in plants and animals that feed upon them (Heaton *et al.* 1986). The herbivores in this study have a 4‰ range of variation, larger than would be expected. This is partly accounted for by the different species considered, but it may also be a reflection of a range of local environmental conditions,

either spatially or temporally, associated with different aridities and local micro-environments. However, the comparison of the results from Temirkash and Tashik show that the mean values for these two sites are similar.

The two dogs have stable isotope results indicating diets at a trophic level intermediate between the herbivores and humans, consistent with the diet expected for domestic dogs.

### *Humans*

In general, the human isotope values suggest a diet largely based on a  $C_3$  terrestrial food chain; however, the results indicate that other food sources with different isotopic values were consumed in varying proportions. The mean differences in  $\delta^{13}C$  between the herbivore and human values (1.0‰ for all individuals, 1.0‰ for Tegiszhol and Temirkash, and 0.8‰ for Tashik) are within the 0–2‰ estimated increase for a trophic level (Bocherens and Drucker 2003). This suggests that the majority of people consumed diets with little or no  $C_4$  input. There are four individuals with notably high  $\delta^{13}C$  values (KZ025  $\delta^{13}C = -16.4‰$ , KZ053  $\delta^{13}C = -15.7‰$ , KZ092  $\delta^{13}C = -16.8‰$  and KZ104  $\delta^{13}C = -16.0‰$ ) suggestive of the addition of  $C_4$  foodstuffs to their diets (although see comments on KZ053 below). Given the faunal data discussed above, it is likely that this  $C_4$  foodstuff was consumed directly by humans, and therefore this probably represents millet consumption and, presumably, cultivation. There are also 11 individuals with intermediate  $\delta^{13}C$  values (between  $-18.0‰$  and  $-17.0‰$ ), although it is not clear if these data represent a small proportion of  $C_4$  foodstuffs in their diets (either through direct consumption or via animals) or if these data represent particularly arid conditions at the time of collagen formation and/or natural variation in human  $\delta^{13}C$  values. On the basis of the human isotopic results presented here, it is not possible to comment on the cultivation and consumption of  $C_3$  plants.

In terms of  $\delta^{15}N$ , even with juveniles removed, the human–animal offset values are higher than that commonly associated with a trophic level enrichment (i.e., 3–5‰; Bocherens and Drucker 2003). The mean human–herbivore offset value of 6.1‰ (5.9‰ at Tegiszhol and Temirkash, and 6.0‰ at Tashik) shows that some of these individuals were consuming some protein from a higher trophic level than the sampled animals. These data strongly indicate that there was a small component of the diet of many individuals with a higher  $\delta^{15}N$  value than the analysed herbivores. The most parsimonious explanation for this is the consumption of a small amount of freshwater fish (or animals that eat freshwater fish). A small number of fish bones have been found at Temirkash through sieving, but the proportion was only 1.5% (unfortunately, these bones were not available for isotopic analysis) and no evidence for fish processing has yet been found through residue analysis of ceramics. Nevertheless, the location of many of the sites along rivers suggests that the inhabitants would have had access to riverine fish. It is possible that fish were treated in such a way that they rarely became part of the archaeological record, such as being prepared and dried or smoked away from the depositional contexts. In general, then, the isotopic evidence indicates that Bronze Age diet in central Kazakhstan was more diverse than traditionally believed. While diet was likely largely based upon domestic herds (note that isotopic analyses cannot distinguish between meat and dairy products), fish formed a notable but varying part of the diet, while some individuals consumed millet.

The statistical analyses above show that this dietary diversity cannot be accounted for by dietary differences between sites. Indeed, the only site with significantly different isotope results was Aschisu, where the proportion of fish consumption was lower than at the other large sites, and fish may not have been consumed at all (particularly by the four individuals with  $\delta^{15}N$  values less

than 12‰). On the other hand, the temporal comparison does account for some of the dietary diversity, in that  $\delta^{13}\text{C}$  values are higher in the Final Bronze Age than the preceding periods. This suggests that the consumption of  $\text{C}_4$  plants, presumably millet, became more important to the diet in this period. It does not necessarily indicate the first consumption of millet in central Kazakhstan but, rather, the first time that it was consumed on a significant scale (cf., Lightfoot *et al.* 2013). Indeed, the presence of carbonized millet remains (see references above) shows that millet was probably present in this Central Asian region in earlier time periods, but infrequently eaten in quantities sufficient to significantly enrich the isotopic signal.

There is one interesting individual, sample KZ053 (Tasyrbai, Fedorovo culture), who has the highest  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values found in this study, while the other individual analysed from this site has more typical values (KZ066;  $\delta^{13}\text{C} = -18.6\text{‰}$  and  $\delta^{15}\text{N} = 11.4\text{‰}$ ). These results may reflect: an aridity effect combined with  $\text{C}_4$  plant consumption (and perhaps migration from a more arid area); a diet very high in freshwater aquatic protein and  $\text{C}_4$  plants; or a diet including large amounts of marine protein (and long-distance migration).

### *Comparison to other studies*

While isotopic research in Central Asia is relatively limited, there are a number of isotopic studies to which the results presented here can be compared. The first isotopic analyses of steppe pastoralists were carried out by O'Connell *et al.* (2003) and suggested that the Eneolithic and Iron Age people sampled consumed significant quantities of fish or other aquatic animals, although the sample size was limited. The study by Svyatko *et al.* (2013) of the Eneolithic to Early Iron Age (2700–1 BC) of the Minusinsk Basin, Russia, also showed that freshwater fish were an important resource throughout this period, and that millet became important in the Late Bronze Age. Murphy *et al.* (2013) have also found isotopic evidence that millet and fish were important dietary components in Early Iron Age southern Siberia. Ventresca Miller and colleagues have used stable isotope analysis and dental analyses (Ventresca Miller *et al.* 2014; Ventresca Miller *et al.* in press) to study diet at Middle Bronze Age Bestamak and Late Bronze Age Lisakovsk in northern Kazakhstan. These analyses suggest that some individuals supplemented their diets with freshwater fish, with little change in diet between the two periods considered. In further support of the consumption of freshwater fish, Privat *et al.* (2007) have used nitrogen and sulphur isotopes to show that freshwater fish were consumed at the Late Bronze Age site of Chicha in Novosibirsk, Russia. The combination here of nitrogen and sulphur isotopic evidence is particularly powerful and convincing. To the east of the study area, Zhang and colleagues (Zhang and Li 2006; Zhang and Zhu 2011; Zhang *et al.* 2009, 2010) have demonstrated that millet consumption was common in Xinjiang, western China, although Bronze Age communities in Xinjiang did not rely on millet as heavily as those in central China.

In summary, although the number of studies is limited, the isotopic data show that fish, and sometimes millet, were part of the diet of steppe 'pastoralists' (who are often assumed to consume mainly terrestrial animal protein), and thus strongly support the interpretations presented here. The true picture of human subsistence in Central Asia is far from clear. However, together with previous studies, this investigation provides a glimpse into the Bronze Age lifestyle in the steppes. The current isotopic evidence suggests that these communities were not as homogeneous in their diets as previously assumed, fitting well with other archaeological evidence for dietary diversity across the steppes (e.g., Frachetti *et al.* 2010; Bendrey 2011; Outram *et al.* 2012; Spengler *et al.* 2014).

## CONCLUSION

The data presented here have shown that the subsistence strategy in Bronze Age central Kazakhstan was not solely focused on terrestrial animals. Instead, fishing and cultivation were notable, if small, parts of the subsistence strategy, with the relative importance of these three dietary components varying for different people and at different times. This dietary diversity can be seen even within sites, indicating that people had real choice in their subsistence strategies. Change can be seen through time, with millet consumption becoming increasingly important in later periods. Through comparison with published research, we have shown that this pattern was not unique to central Kazakhstan—the, admittedly limited, isotopic and archaeobotanical evidence indicates the importance of these resources at various times and locations cross the steppes.

The importance of fishing and cultivation requires the economic basis and daily lives of these communities to be reconsidered. Fishing and cultivation would have required considerable inputs of time, energy and resources on a (semi-)regular basis. This has important implications for our understanding of what it means to be a pastoralist. We suggest that the use of domestic herds does not preclude the procurement of other food resources—millet, for example, requires relatively little human input between sowing and harvesting (Rachie 1975; Rao 1989), and the evidence presented here and elsewhere shows that these pastoralist communities were responding in diverse ways to the opportunities presented by the world around them.

## ACKNOWLEDGEMENTS

The authors are grateful to Catherine Kneale, Mike Hall and James Rolfe (University of Cambridge) for their assistance with the isotopic analysis. We also wish to thank Professor Beisenov for providing the samples from Kyzyl. We are grateful to the European Research Council and Darwin College, Cambridge (EL) for financial support. GM-M's postdoctoral fellowship is being funded by the European Union Structural Funds project 'Postdoctoral Fellowship Implementation in Lithuania'.

## REFERENCES

- Ambrose, S. H., 1990, Preparation and characterization of bone and tooth collagen for isotopic analysis, *Journal of Archaeological Science*, **17**(4), 431–51.
- Ambrose, S. H., and Norr, L., 1993, Experimental evidence for the relationship of the carbon isotope ratios of whole diet and dietary protein to those of bone collagen and carbonate, in *Prehistoric human bone: archaeology at the molecular level* (eds. J. Lambert and G. Grupe), Springer-Verlag, New York.
- Anthony, D., 2007, *The horse, the wheel, and language: how Bronze-Age riders from the Eurasian steppes shaped the modern world*, Princeton University Press, Princeton, NJ.
- Beisenov, A., and Varfolomeev, V., 2008, *Могильник Бегазы. Центральный Казахстан в Бегазы–Дандыбаевскую эпоху (Begazy cemetery. Central Kazakhstan in the Begazy–Dandybai era)*, Almaty.
- Bendrey, R., 2011, Some like it hot: Environmental determinism and the pastoral economies of the later prehistoric Eurasian steppe, *Pastoralism: Research, Policy and Practice*, **1**(8), 1–16.
- Bocherens, H., and Drucker, D., 2003, Trophic level isotopic enrichment of carbon and nitrogen in bone collagen: case studies from recent and ancient terrestrial ecosystems, *International Journal of Osteoarchaeology*, **13**(1–2), 46–53.
- Bogaard, A., Heaton, T. H. E., Poulton, P., and Merbach, I., 2007, The impact of manuring on nitrogen isotope ratios in cereals: archaeological implications for reconstruction of diet and crop management practices, *Journal of Archaeological Science*, **34**(3), 335–43.
- Chang, C., Benecke, N., Grigoriev, F. P., Rosen, A. M., and Tourtellotte, P. A., 2003, Iron Age society and chronology in south-east Kazakhstan, *Antiquity*, **77**(296), 298–312.
- De Niro, M. J., 1985, Postmortem preservation and alteration of *in vivo* bone collagen isotope ratios in relation to paleodietary reconstruction, *Nature*, **317**(6040), 806–9.

- Farquhar, G. D., Ehleringer, J. R., and Hubick, K. T., 1989, Carbon isotope discrimination and photosynthesis, *Annual Review of Plant Physiology and Plant Molecular Biology*, **40**, 503–37.
- Frachetti, M., 2002, Bronze Age exploitation and political dynamics of the eastern Eurasian steppe zone, in *Ancient interactions: East and West in Eurasia* (eds. K. Boyle, C. Renfrew and M. Levine), 161–70, McDonald Institute for Archaeological Research, Cambridge.
- Frachetti, M. D., 2006, The Dzhungar Mountain Archaeology Project: reconstructing Bronze Age life in the mountains of eastern Kazakhstan, in *Beyond the steppe and the sown* (eds. D. L. Peterson, L. M. Popova and A. T. Smith), 122–41, Brill, Boston.
- Frachetti, M. D., Spengler, R. N., Fritz, G. J., and Mar'yashev, A. N., 2010, Earliest direct evidence for broomcorn millet and wheat in the central Eurasian steppe region, *Antiquity*, **84**, 993–1010.
- Fuller, B. T., Fuller, J. L., Harris, D. A., and Hedges, R. E. M., 2006, Detection of breastfeeding and weaning in modern human infants with carbon and nitrogen stable isotope ratios, *American Journal of Physical Anthropology*, **129**, 279–93.
- Gayduchenko, L. L., 2002, Organic remains from fortified settlements and necropolis of the 'Country of Towns', In *Complex societies of central Eurasia from the 3rd to the 1st millennium BC: regional specifics in light of global models*, vol. 2: *The Iron Age; archaeoecology, geoarchaeology, and palaeogeography; beyond central Eurasia* (eds. K. Jones-Bley and D. G. Zdanovich), 400–18, Institute of Man, Washington, DC.
- Gimbutas, M., 1973, Old Europe c.7000–3500 B.C.: the earliest European civilization before the infiltration of the Indo-European peoples, *Journal of Indo-European Studies*, **1**, 1–20.
- Hanks, B., 2010, Archaeology of the Eurasian steppes and Mongolia, *Annual Review of Anthropology*, **39**, 469–86.
- Hanks, B. K., Epimakhov, A. V., and Renfrew, A. C., 2007, Towards a refined chronology for the Bronze Age of the southern Urals, Russia, *Antiquity*, **81**, 353–67.
- Hartman, G., 2012, Impacts of environmental deterioration on the carbon isotope values of modern vegetation and gazelles in the southern Levant: predicting the severity of the Younger Dryas, *Palaeogeography, Palaeoclimatology, Palaeoecology*, **321–2**, 55–64.
- Heaton, T. H. E., Vogel, J. C., Vonlachevallerie, G., and Collett, G., 1986, Climatic influence on the isotopic composition of bone nitrogen, *Nature*, **322**(6082), 822–3.
- Hedges, R. E. M., 2003, On bone collagen—apatite—carbonate isotopic relationships, *International Journal of Osteoarchaeology*, **13**(1–2), 66–79.
- Hedges, R. E. M., and Reynard, L., 2007, Nitrogen isotopes and the trophic level of humans in archaeology, *Journal of Archaeological Science*, **34**, 1240–51.
- Hedges, R. E. M., Clement, J. G., Thomas, D. L., and O'Connell, T. C., 2007, Collagen turnover in the adult femoral mid-shaft: modelled from anthropogenic radiocarbon tracer measurements, *American Journal of Physical Anthropology*, **133**, 808–16.
- Hoefs, J., 2004, *Stable isotope geochemistry*, Springer Verlag, Heidelberg.
- Howland, M. R., Corr, L. T., Young, S. M. M., Jones, V., Jim, S., Van Der Merwe, N. J., Mitchell, A. D., and Evershed, R. P., 2003, Expression of the dietary isotope signal in the compound-specific delta(13) values of pig bone lipids and amino acids, *International Journal of Osteoarchaeology*, **13**(1–2), 54–65.
- Jim, S., Jones, V., Ambrose, S. H., and Evershed, R. P., 2006, Quantifying dietary macronutrient sources of carbon for bone collagen biosynthesis using natural abundance stable carbon isotope analysis, *British Journal of Nutrition*, **95**, 1055–62.
- Katzenberg, M. A., and Lovell, N. C., 1999, Stable isotope variation in pathological bone, *International Journal of Osteoarchaeology*, **9**(5), 316–24.
- Koryakova, L., and Epimakhov, A. V., 2007, *The Urals and western Siberia in the Bronze and Iron Ages*, Cambridge University Press, Cambridge.
- Kukushkin, I. A., 2006, Могильник Аяпберген—раннеандроновский памятник центрального Казахстана (Ayarbergen cemetery—early Andronovo site in central Kazakhstan), in *Изучение памятников археологии Павлодарского Прииртышья* (Study of archaeological sites of Pavlodar region) (ed. V. K. Merc), 50–69, Pavlodar.
- Kukushkin, I. A., 2007, Археологические исследования могильника Ащису. Курган 1 (Archaeological research repository Aschisu. Mound 1), in *Историко-культурное наследие Сарыарки* (Historical and cultural heritage of Saryaka) (ed. V. G. Loman), 40–64, Karaganda.
- Kukushkin, I. A., 2010, Новые исследования на могильнике Ащису (New research on the Aschisu cemetery), *Вестник археологии, антропологии и этнографии* (*Journal of Archaeology, Anthropology and Ethnography*), **1**(12), 71–5.

- Kukushkin, I. A., 2012, Могильник Ащису. Курган с металлическим сосудом (Aschisu cemetery. Mound with a metal vessel), in *Археология и история Сарыарки (Archaeology and History of Saryarka)* (ed. R. M. Zhumashev), 63–81, Karaganda.
- Kukushkin, I. A., and Loman, V. G., 2004, Курган Жартаc (Zhartas kurgan), *Известия Национальной академии наук Республики Казахстан. Серия общественных наук (Proceedings of the National Academy of Sciences of the Republic of Kazakhstan. Social Sciences Series)*, **1**(247), S. 6367.
- Kukushkin, I. A., and Loman, V. G., 2011, Курган гунно-сарматского времени на могильнике Ащису (Hun-Sarmatian date for Aschisu burial), *Вопросы археологии Казахстана (Questions in Kazakh Archaeology)*, **3**, 315–24.
- Kuzmina, E. E., 2003, Origins of pastoralism in the Eurasian steppes, in *Prehistoric steppe adaptation and the horse* (eds. M. A. Levine, A. C. Renfrew and K. Boyle), 203–32, McDonald Institute for Archaeological Research, Cambridge.
- Lightfoot, E., Liu, X., and Jones, M. K., 2013, Why move starchy cereals? A review of the isotopic evidence for prehistoric millet consumption across Eurasia, *World Archaeology*, **45**(4), 574–623.
- Loman, V. G., and Kukushkin, I. A., 2009, Могильник Кудрявая Сопка-1 (Kudryavaya Sopka-1 cemetery), in *Изучение историко культурного наследия Центральной Евразии (Study of the historical and cultural heritage of Central Eurasia)* (ed. V. V. Varfolomeev), 91–6, Karaganda.
- Loman, V. G., and Kukushkin, I. A., 2011, К вопроу о культуре плодородия у андроновцев (федоровцев) (On the fertility cult in the Andronovo (Fedorovo)), in *Материалы международной научной конференции «Археология Казахстана в эпоху независимости: итоги, перспективы», посвященной 20-летию независимости Песпублики Казахстан и 20-летию Института археологии им. А. Х. Маргулана (Proceedings of the international scientific conference 'Archaeology of Kazakhstan independence era: results and prospects', marking the 20th anniversary of Kazakhstan's independence and the 20th anniversary of the Institute of Archaeology named after A. H. Margulan)*, vol. I, 278–80, Almaty.
- Loman, V. G., and Kukushkin, I. A., 2012, Могильник Дарьинский (Darya cemetery), in *Археология и история Сарыарки (Archaeology and history of Saryarka)* (ed. R. M. Zhumashev), 81–106, Karaganda.
- Mekota, A. M., Grupe, G., Ufer, S., and Cuntz, U., 2006, Serial analysis of stable nitrogen and carbon isotopes in hair: monitoring starvation and recovery phases of patients suffering from anorexia nervosa, *Rapid Communications in Mass Spectrometry*, **20**(10), 1604–10.
- Murphy, E. M., Schulting, R., Beer, N., Chistov, Y., Kasparov, A., and Pshenitsyna, M., 2013, Iron Age pastoral nomadism and agriculture in the eastern Eurasian steppe: implications from dental palaeopathology and stable carbon and nitrogen isotopes, *Journal of Archaeological Science*, **40**(5), 2547–60.
- O'Connell, T. C., and Hedges, R. E. M., 1999, Investigations into the effect of diet on modern human hair isotopic values, *American Journal of Physical Anthropology*, **108**(4), 409–25.
- O'Connell, T. C., Levine, M. A., and Hedges, R. E. M., 2003, The importance of fish in the diet of central Eurasian peoples from the Mesolithic to the Early Iron Age, in *Prehistoric steppe adaptation and the horse* (eds. M. A. Levine, A. C. Renfrew and K. Boyle), 253–68, McDonald Institute for Archaeological Research, Cambridge.
- O'Leary, M. H., 1988, Carbon isotopes in photosynthesis, *Bioscience*, **38**(5), 328–35.
- Outram, A. K., Kasparov, A., Stear, N. A., Varfolomeev, V., Usmanova, E., and Evershed, R. P., 2012, Patterns of pastoralism in later Bronze Age Kazakhstan: new evidence from faunal and lipid residue analyses, *Journal of Archaeological Science*, **39**, 2424–35.
- Outram, A. K., Stear, N. A., Kasparov, A., Usmanova, E., Varfolomeev, V., and Evershed, R. P., 2011, Horses for the dead: funerary foodways in Bronze Age Kazakhstan, *Antiquity*, **85**, 116–28.
- Privat, K. L., O'Connell, T. C., and Hedges, R. E. M., 2007, The distinction between freshwater- and terrestrial-based diets: methodological concerns and archaeological applications of stable sulphur isotope analysis, *Journal of Archaeological Science*, **34**(8), 1197–204.
- Privat, K. L., Schneeweß, J., Benecke, N., Vasil'ev, S. K., O'Connell, T. C., Hedges, R. E. M., and Craig, O., 2005, Economy and diet at the Late Bronze Age – Iron Age site of Čiča: artefactual, archaeozoological and biochemical analyses, *Eurasia Antiqua*, **11**, 419–48.
- Pyanokov, V. I., Gunin, P. D., Tsoog, S., and Black, C. C., 2000, C<sub>4</sub> plants in the vegetation of Mongolia: their natural occurrence and geographical distribution in relation to climate, *Oecologia*, **123**(1), 15–31.
- Rachie, K. O., 1975, *The millets: importance, utilisation and outlook*, International Crops Research Institute for the Semi-arid Tropics, Hyderabad.
- Rao, M. V., 1989, Inaugural address, in *Small millets in global agriculture* (eds. A. Seetharam, K. W. Riley and G. Harinarayana), Oxford & IBH Publishing, New Delhi.
- Renfrew, C., 1987, *Archaeology and language: the puzzle of Indo-European origins*, Jonathan Cape, London.



- Schoeninger, M. J., and De Niro, M. J., 1984, Nitrogen and carbon isotopic composition of bone-collagen from marine and terrestrial animals, *Geochimica et Cosmochimica Acta*, **48**(4), 625–39.
- Spengler, R. N., Frachetti, M. D., and Fritz, G. J., 2013, Ecotopes and herd foraging practices in the steppe/mountain ecotone of Central Asia during the Bronze and Iron Ages, *Journal of Ethnobiology*, **33**(1), 125–47.
- Spengler, R. N., Frachetti, M. D., Doumani, P., Rouse, L., Cerasetti, B., Bullion, E., and Mar'yashev, A., 2014, Early agriculture and crop transmission among Bronze Age mobile pastoralists of Central Asia, *Proceedings of the Royal Society Series B*, **281**(1783) (available online).
- Stevens, R. E., and Hedges, R. E. M., 2004, Carbon and nitrogen stable isotope analysis of northwest European horse bone and tooth collagen, 40,000 BP–present: palaeoclimatic interpretations, *Quaternary Science Reviews*, **23**(7–8), 977–91.
- Svyatko, S. V., Mallory, J. P., Murphy, E. M., Polyakov, A. V., Reimer, P. J., and Schulting, R. J., 2009, New radiocarbon dates and a review of the chronology of prehistoric populations from the Minusinsk Basin, southern Siberia, Russia, *Radiocarbon*, **51**(1), 243–73.
- Svyatko, S. V., Schulting, R. J., Mallory, J., Murphy, E. M., Reimer, P. J., Khartanovich, V. I., Chistov, Y. K., and Sablin, M. V., 2013, Stable isotope dietary analysis of prehistoric populations from the Minusinsk Basin, southern Siberia, Russia: a new chronological framework for the introduction of millet to the eastern Eurasian steppe, *Journal of Archaeological Science*, **40**, 3936–45.
- Tieszen, L. L., and Fagre, T., 1993, Effect of diet quality on the isotopic composition of respiratory CO<sub>2</sub>, bone collagen, bioapatite and soft tissues, in *Prehistoric human bone: archaeology at the molecular level* (eds. J. B. Lambert and G. Grupe), Springer-Verlag, Berlin.
- Toderich, K., Black, C. C., Juylova, E., Kozan, O., Mukimov, T., and Matsuo, N., 2004, C3/C4 plants in the vegetation of Central Asia, geographical distribution and environmental adaptation in relation to climate, in *Climate change and terrestrial carbon sequestration in Central Asia* (eds. R. Lal, M. Suleimenov, B. A. Stewart, D. O. Hansen and P. Doraiswamy), 33–63, Taylor and Francis, London.
- Varfolomeev, V., 2007a, Погребения культуры валиковой керамики в урочище Темиркаш (Burial culture of roll ceramics in the area of Temirkash), in *Кадырбаевские чтения. Материалы международной научной конференции (Kadyrbaevskie reading. Materials of international conference)*, 50–7, Aktobe.
- Varfolomeev, V., 2007b, Могилиник Тегисжол по раскопкам 2005 г. (Tegizhol Cemetery Excavation 2005), in *Историко-культурное наследие Сарыарки (Historical and cultural heritage of Saryarka)* (ed. V. G. Loman), 65–84, Karaganda.
- Varfolomeev, V., 2011a, Кент—город бронзового века. Новые исследования в эпоху независимости (Kent—a Bronze Age city. New research in the era of independence), in *Свидетели тысячелетий: археологическая наука Казахстана за 20 лет (1991–2011) (Witnesses of millennia: archaeological science in Kazakhstan for 20 years (1991–2011))*, 85–97, Institute of Archaeology A. N. Margulan, Almaty.
- Varfolomeev, V., 2011b, Тасмолинское погребение с наборным поясом из могилиника Тегисжол (The Tasmola burial in which a girdle from the Tegizhol burial ground was found), *Вопросы археологии Казахстана (Questions in Kazakh Archaeology)*, **3**, 315–24.
- Varfolomeev, V., 2013, Die Begazy–Dandybaj-Kultur (The Begazy–Dandybaj culture), in *Unbekanntes Kasachstan. Archäologie im Herzen Asiens (Unknown Kazakhstan. Archaeology in the heart of Asia)*, 483–98, German Mining Museum, Bochum.
- Ventresca Miller, A., Usmanova, E., Logvin, V., Kalieva, S., Shevnina, I., Logvin, A., Kolbina, A. and Suslov, A., in press, Dental health, diet, and social transformations in the Bronze Age: comparative analysis of pastoral populations in northern Kazakhstan, *Quaternary International*.
- Ventresca Miller, A., Usmanova, E., Logvin, V., Kalieva, S., Shevnina, I., Logvin, A., Kolbina, A., Suslov, A., Privat, K., Haas, K., and Rosenmeier, M., 2014, Subsistence and social change in central Eurasia: stable isotope analysis of populations spanning the Bronze Age transition, *Journal of Archaeological Science*, **42**, 525–38.
- Vogel, J. C., and Van Der Merwe, N. J., 1977, Isotopic evidence for early maize cultivation in New York State, *American Antiquity*, **42**(2), 238–42.
- Zhang, Q., and Li, S., 2006, Xinjiang Nileke Xian Kongkeke yihao mudi gudai jumin de shiwu jiegou fenxi (Analysis on paleodiet of ancient inhabitants in no. 1 cemetery of Qiongkeke at Nika County, Xinjiang), *Xiyu Yanjiu (The Western Regions Studies)*, **4**, 78–82.
- Zhang, Q., and Zhu, H., 2011, Xinjiang Gumugou mudi rengu de wending tongweisu fenxi: zaoqi Luobubo xianmin yinshi jiegou chutan (Stable isotopic analysis on human bones from Gumugou cemetery, Xinjiang: paleodietary reconstruction on Luobubo population), *Xiyu Yanjiu (The Western Regions Studies)*, **3**, 91–6.
- Zhang, Q., Chang, X., and Liu, G., 2009, Xinjiang Balikun xian Heigouliang mudi chutu rengu de shixing fenxi (Stable isotopic analysis on human bones from Heigouliang cemetery in Barkol, Xinjiang), *Xiyu Yanjiu (The Western Regions Studies)*, **3**, 45–9.

Zhang, Q., Chang, X., and Liu, G., 2010, Xinjiang Hami Tianshanbeilu mudi chutu rengu de wending tongweisu fenxi (Stable isotopic analysis of human bones unearthed from Tianshan Beilu cemetery in Hami, Xinjiang), *Xiyu Yanjiu (The Western Regions Studies)*, **2**, 38–43.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

Appendix 1.

Appendix 2.