

LETTERS
TO THE EDITOR

Stereoselective Carbocyclization of Hanphyllin
with *N*-Bromosuccinimide

N. Merkhatuly, S. K. Zhokizhanova, L. T. Balmagambetova, and S. M. Adekenov

Buketov Karaganda State University,
ul. Universitetskaya 28, Karaganda, 100028 Kazakhstan
e-mail: galija@kargu.krg.kz

Received February 20, 2006


DOI: 10.1134/S1070363206080378

It is known that sesquiterpene γ -lactones of the germacrane structure (germacranolides) in acid media undergo facile intramolecular cyclization to form bicyclic (eudesmane, guaiane, etc.) sesquiterpenoids [1–3].

We performed a stereoselective transannular carbocyclization of 1(10)*E*,4*E*-germacranolide hanphyll-

lin (**I**) with *N*-bromosuccinimide to obtain novel *trans*-fused (5 α -*H*) eudesmane sesquiterpene γ -lactones **II** and **III** in 30 and 40% yields, respectively.

The stereoselectivity of the reaction is probably explained by the *chair*–*chair* conformation of hanphyllin (**I**) [4].



3 β -Hydroxy-5,7 α -*H*-eudesma-1(2),4(15),11(13)-trien-6,12-olide (II) and 3 β ,4 α -dihydroxy-5,7 α -*H*-eudesma-1(2),11(13)-dien-6,12-olide (III). To a solution of 2.2 g of hanphyllin in 5 ml of 5% aqueous acetone, 0.142 g of finely ground *N*-bromosuccinimide was added with stirring. The mixture was stirred for 1.5 h and then, at 25–30°C, poured into 3% HCl. The reaction products were extracted with ethyl acetate, the extract was dried with MgSO₄, the solvent was removed by vacuum distillation, and the residue was subjected to column chromatography on silica gel, eluent hexane–ethyl acetate.

Compound **II**, yield 0.059 g (30%), colorless crystals, mp 142–144°C. IR spectrum, ν , cm⁻¹: 3450 (HO),

1750 (C=O), 1650, 1640 (C=C). ¹H NMR spectrum, δ , ppm: 0.90 s (3H, CH₃-C¹⁰), 5.28 d (1H, HC¹, *J*_{HH} 11.0 Hz), 5.40 q (1H, HC², *J*_{HH} 11.0, 9.0 Hz), 4.40 br.d (1H, HC³, *J*_{HH} 9.0 Hz), 2.24 br.d (1H, HC⁵, *J*_{HH} 11.0 Hz), 3.98 t (1H, HC⁶, *J*_{HH} 11.0 Hz), 5.40 d (1H, HC¹³, *J*_{HH} 3.0 Hz), 6.06 d (1H, HC¹³, *J*_{HH} 3.0 Hz), 5.25 d (1H, HC¹⁵, *J*_{HH} 1.0 Hz), 5.90 d (1H, HC¹⁵, *J*_{HH} 1.0 Hz). Found, %: C 72.97; H 7.21. C₁₅H₁₈O₃. Calculated, %: C 73.17; H 7.31.

Compound **III**, yield 0.084 g (40%), colorless crystals, mp 161–163°C. IR spectrum, ν , cm⁻¹: 3450 (OH), 1750 (C=O), 1650 (C=C). ¹H NMR spectrum, δ , ppm: 1.26 s (3H, CH₃-C⁴), 0.94 s (3H, CH₃-C¹⁰), 5.29 d (1H, HC¹, *J*_{HH} 11.0 Hz), 5.41 q (1H, HC², *J*_{HH}

11.0, 9.0 Hz), 4.45 d (1H, HC³, J_{HH} 9.0 Hz), 2.18 d (1H, HC⁵, J_{HH} 11.0 Hz), 6.14 d (1H, HC¹³, J_{HH} 3.0 Hz). Mass spectrum, m/z (I_{rel} , %): 265 (44.7) [$M + H$], 247 (61.8), 207 (100), 147 (28.9), 119 (19.7), 81 (25.0), 55 (25.0), 43 (69.7). Found, %: C 72.99; H 7.38. C₁₅H₂₀O₄. Calculated, %: C 73.17; H 7.57.

The IR spectra were measured on an Avatar-360 instrument in KBr. The ¹H NMR spectra were taken on a Bruker WP-200SY instrument (200.13 MHz) in CDCl₃, internal reference TMS. The mass spectrum

was taken on a Finnigan MAT-8200 instrument, ionizing energy 70 eV.

REFERENCES

1. Jain, T.C. and McCloskey, J.E., *Tetrahedron*, 1975, vol. 31, no. 9, p. 2211.
2. Parodi, F.J., Fronscek, F.R., and Fischer, H.N., *J. Nat. Prod.*, 1989, vol. 52, no. 3, p. 423.
3. Barrero, A.F., Oltra, I.E., Morales, V., and Alvarez, M.A., *J. Nat. Prod.*, 1997, vol. 60, no. 10, p. 1034.
4. Tashkhodzhaev, B. and Abduazimov, B.Kh., *Khim. Prirodn. Soedin.*, 1997, no. 2, p. 497.